

MET Block & Variables

- ❑ Changes to missingET package
- ❑ Root-tuple blocks from met_analyze
- ❑ Missing transverse energy variables
- ❑ Visible and scalar energy variables
- ❑ MET example plots
- ❑ What's next?

Alan L. Stone - Louisiana Tech University

27 February 2002
DO Jet/MET Mtg.



Alan L. Stone
Louisiana Tech University




Changes to missingET package

- ❑ New algorithms to calculate new variables
- ❑ Removed specialized constructor used only by PMCS
 - Use default constructor with set methods for some variables
- ❑ Removed boolean absET2
 - Did nothing & original purpose was not clear
- ❑ Added rcps for:
 - cellThreshold (>100 MeV)
 - towerThreshold (>200 MeV)
 - etaLimit ($\eta < 33$)
- ❑ Initialize function:
 - `void setAllZero (void);`
- ❑ All variables placed in thumbnail by default
 - Do we want this?
 - Increased packing (S.B.)
- ❑ Full compliment of *set/get/print* methods for all variables
 - old methods were left unchanged




Algorithm Details

$$E_{x,y}^{Vis.CAL} = \sum_{cells} E_i^{x,y}$$



$$ME_{x,y}^{CAL} = -E_{x,y}^{Vis.CAL}$$




$$MET^{CAL} = \sqrt{(ME_x^{CAL})^2 + (ME_y^{CAL})^2}$$



$$SET^{CAL} = \sum_{cells} \sqrt{(E_i^x)^2 + (E_i^y)^2}$$

$$ME_{x,y}^{CAL+MUON} = ME_{x,y}^{CAL} - P_{x,y}^{MUON,vis}$$

$$P_{x,y}^{Muon,vis} = \sum_{muons} p_i^{x,y}$$



$$MET^{CAL+MUON} = \sqrt{(ME_x^{C+M})^2 + (ME_y^{C+M})^2}$$

MET

met_analyze package

Missing Transverse Energy: *ScalarET, METx, METy, MET*

- Calorimeter + ICD → cells or towers
- Calorimeter + ICD with muon correction → cells or towers
 - (tight muons in *muonid*)

Visible Energy: *SET, VETx, VETy, VET*

- Calorimeter + ICD → cells or towers
 - Above/Below ieta limit (32 inclusive)
 - Signal/Noise ($E_{\text{cells}} > 100 \text{ MeV}$, $E_{\text{towers}} > 200 \text{ MeV}$)
 - ICD cells only; NADA killed cells
- Muon: *Px, Py, Pz, Pt*

METMET

Old variables (jetanalyze METOLD block)

MET, MEx, MEy, SET, MEphi
METNE, MExNE, MEyNE, SETNE, MENEphi
METWE, MExWE, MEyWE, SETWE, MEWEphi
SMuX, SMuY, SMuZ, Zvtx

METRING

Ring variables for revertexing (EM & Had. components, 42 ieta segments)

Nrings
RingEMx[42], RingEMy[42]
RingHDx[42], RingHDy[42]



METOLD block in **jetanalyze** (METMET in **met_analyze**)


→ Calculated from towers in CalData chunk

MET Missing Et Calorimeter only
METx x component of MET - Cal only (can be negative)
METy y component of MET - Cal only (can be negative)
MEphi METx/METy
SET Scalar ET

METNE Same as MET with threshold of 200 MeV/tower
METNEx Same as METx w/threshold of 200 MeV/tower
METNEy Same as METy w/threshold of 200 MeV/tower
MEphiNE METNEx/METNEy
SETNE Same as SET w/threshold of 200 MeV/tower

METWE Same as METNE with $|i\eta| < 33$
METWEx Same as METNEx with $|i\eta| < 33$
METWEy Same as METNEy with $|i\eta| < 33$
MEphiWE METWEx/METWEy
SETWE Same as SETNE with $|i\eta| < 33$

Zvtx primary reconstructed Z position of vertex
SMuX px sum of the "good muons"
SMuY py sum of the "good muons"
SMuPt Transverse momentum of the "good muons"



Date: Mon, 19 Nov 2001 23:00:25 +0100 (CET)
From: Gregorio Bernardi <gregorio@in2p3.fr>
To: d0jetmet@fnal.gov
Subject: Upgrade of the MET Block/scheme.

SET: scalar ET obtained as the sum of cell energy*abs(sin(theta)). A cell with negative energy will give a NEGATIVE contribution to SET.

VETx: x of vect. ET obtained as the sum of the cell energy*sin(th)*cos(ph)

VETy: y of vect. ET obtained as the sum of the cell energy*sin(th)*sin(ph)

VETz: z of vect. ET obtained as the sum of the cell energy*cos(th)

(in these 3 variables the energy can be positive or negative)

$$VET = \sqrt{VETx^2 + VETy^2}$$

The x and y component of the Missing ET (METx, METy) are simply:

$$METx = -VETx$$

$$METy = -VETy$$

The missing transverse energy is obtained by $MET = \sqrt{METx^2 + METy^2}$, and of course $MET = VET$.

In the detailed variables, we use VETx, VETy, VETz in order to have a symmetric (same sign) treatment of calorimeter and muons, e.g.


$$VETx = +VETCALOx + VETMUONx, \text{ etc...}$$



Structure of the future MET block

Global variables:

SETT	Scalar Et constructed from Cal-ICD Towers	(=+SETTAS +SETTBS)
METT _x	x component constructed from Cal-ICD Towers	(=-VETTAS _x -VETTBS _x)
METT _y	y component constructed from Cal-ICD Towers	(=-VETTAS _y -VETTBS _y)
METT	Missing Et constructed from Cal-ICD Towers	
METTM	Scalar Et constructed from Cal-ICD Towers&Muon	(+SETT +SETMUON)
METTM _x	x component constructed from Cal-ICD Towers&Muon	(-VETT _x -VETMUON _x)
METTM _y	y component constructed from Cal-ICD Towers&Muon	(-VETT _y -VETMUON _y)
METTM	Missing Et constructed from Cal-ICD Towers&Muon	
SETC	Scalar Et constructed from Cal-ICD Cells	(+SETCAS +SETCBS)
METC _x	x component constructed from Cal-ICD Cells	(-VETCAS _x -VETCBS _x)
METC _y	y component constructed from Cal-ICD Cells	(-VETCAS _y -VETCBS _y)
METC	Missing Et constructed from Cal-ICD Cells	
SETCM	Scalar Et constructed from Cal-ICD Cells&Muon	(+SETC +SETMUON)
METCM _x	x component constructed from Cal-ICD Cells&Muon	(-VETC _x -VETMUON _x)
METCM _y	y component constructed from Cal-ICD Cells&Muon	(-VETC _y -VETMUON _y)
METCM	Missing Et constructed from Cal-ICD Cells&Muon	



SETTAS	Scalar sum of Towers Above eta limit, $E_{tow} > twr-thresh$.
VETTASx	Vectorial sum of Towers Above eta limit, $E_{tow} > twr-thresh$.
SETTBS	Scalar sum of Towers Below eta limit, $E_{tow} > twr-thresh$.
VETTBSx	Vectorial sum of Towers Below eta limit, $E_{tow} > twr-thresh$.
SETTAN	Scalar sum of Towers Above eta limit, $E_{tow} < twr-thresh$.
VETTANx	Vectorial sum of Towers Above eta limit, $E_{tow} < twr-thresh$.
SETTBN	Scalar sum of Towers Below eta limit, $E_{tow} < twr-thresh$.
VETTBNx	Vectorial sum of Towers Below eta limit, $E_{tow} < twr-thresh$.

SETCAS	Scalar sum of Cells Above eta limit, $E_{cell} > cel-thresh$.
VETCASx	Vectorial sum of Cells Above eta limit, $E_{cell} > cel-thresh$.
SETCBS	Scalar sum of Cells Below eta limit, $E_{cell} > cel-thresh$.
VETCBSx	Vectorial sum of Cells Below eta limit, $E_{cell} > cel-thresh$.
SETCAN	Scalar sum of Cells Above eta limit, $E_{cell} < cel-thresh$.
VETCANx	Vectorial sum of Cells Above eta limit, $E_{cell} < cel-thresh$.
SETCBN	Scalar sum of Cells Below eta limit, $E_{cell} < cel-thresh$.
VETCBNx	Vectorial sum of Cells Below eta limit, $E_{cell} < cel-thresh$.

T
O
W
E
R
S

C
E
L
L
S



SETICD Scalar sum of ICD cells only
VETICDx Vectorial sum of ICD cells only

SETNADAx Scalar sum of NADA cells
VETNADAx Vectorial sum of NADA cells


SETMUONx Scalar sum of MUONs
VETMUONx Vectorial sum of MUONs

The **MUON**, **ICD**, **NADA** quantities are given for systematic studies. They are not used explicitly in the global MET/SET calculations done in this block, since the ICD is already included in the Tower and Cells variables, and since NADA hot cells have already been suppressed. However, if NADA is ran in shadow mode then the NADA quantities can be used to obtain a NADA corrected missing/Scalar ET. Conversely, if NADA has been run in killing mode, the non-NADA corrected MET can be trivially restored using the detailed NADA variables.


The revertexing block corresponds to the **VETTAS+VETTBS** quantities.
(i.e. the total vectorial E_T obtained from towers above tower threshold).

The WE/NE concept of the old MET block is implemented in this scheme by the 2 complementary quantities:


SETTAS (towers above eta limits) and **SETTBS** (towers below eta limit),
i.e. **SETT=SETTAS+SETTBS** (similary for VETxy, MET).




New MET Root-tuple Block



Missing ET, cells - reco_analyze



Missing ET, cells - Muon corrected




27 February 2002
DO Jet/MET Mtg.



Alan L. Stone
Louisiana Tech University



Missing ET, towers - Muon corrected




27 February 2002
DO Jet/MET Mtg.




Alan L. Stone
Louisiana Tech University



Visible ET, cells: Signal vs Noise



Visible ET, towers: Below/Above eta limit



What's Next?

- ❑ Thoroughly test new variables
 - Need everyone to revisit their favorite data & macros
 - Can one reproduce plots made with METOLD with variables in MET (and METMET)?
 - Check default limits in RCP (energy, ieta)
 - Revertexing with rings (*Joe Steele*)
 - Still need NADA (killed cells)
- ❑ Strip METOLD block from jetanalyze and METMET block met_analyze
 - Which one first? Is everyone convinced that METOLD variables are just a subset of the new MET variables?
- ❑ Should we only use tight muons to correct the calorimeter missing ET?
 - What is the muon ID group doing?
- ❑ Going into t02.06.00 & p11...
 - missingET v00-03-01 & met_analyze v00-03-00

